

Vitamin D: An Absolute Requirement for Healthy Living

Everyone should have his or her vitamin D level checked at least once a year (infants through the elderly).

Below are conditions that have been associated with vitamin D levels:

< 10 ng/mL	Severely deficient
< 15 ng/mL	Risk of rickets ⁱ
< 20 ng/mL	75% greater risk of colon cancer ⁱⁱ
< 30 ng/mL	Deficient
	Increased calcium loss from bones, osteoporosis ⁱⁱⁱ
	Poor wound healing ^{iv}
	Increased muscle pain ^v
	Increased joint and back pain ^{vi}
	Greater risk of depression ^{vii}
	Increased diabetes ^{viii}
	Increased schizophrenia ^{ix}
	Increased migraines ^x
	Increased autoimmune disease (lupus, scleroderma) ^{xi}
	Increased allergies
	Increased preeclampsia ^{xii}
	Increased inflammation
30–50 ng/mL	Suboptimal levels
< 34 ng/mL	Twice the risk of heart attack ^{xiii}
< 36 ng/mL	Increased incidence of high blood pressure ^{xiv}
< 40 ng/mL	Three times the risk of multiple sclerosis ^{xv}
50–80 ng/mL	Optimal levels
> 50 ng/mL	50% reduction in breast cancer, decreased risk of all solid cancers ^{xvi}
80–100 ng/mL	Slowing of cancer growth in patients with cancer ^{xvii}
> 100 ng/mL	Increased risk of toxic symptoms (hypercalcemia)^{xviii}

Natural Production of Vitamin D

Your skin makes vitamin D when it is exposed to a pinkish dose of sunlight. How much vitamin D you make depends on your age, how much skin is uncovered, and your skin tone. Without sunblock and with arms and legs exposed, your skin will make 10,000 to 15,000 units of vitamin D in one pinkish sun exposure, on average. Sunblock with an SPF of more than 15 blocks 100% of vitamin D production in the skin.

Depending on the latitude at which you live, you may only get enough radiation from the sun for vitamin D production between May and October. Also, the darker your skin, the more sun you need to make enough vitamin D.

Vitamin D Supplementation Doses

Normal dosing of vitamin D depends on your blood levels. Treatment doses for blood level ranges are:

<10 ng/mL –	10,000 units per day
10–20 ng/mL –	10,000 units per day
20–30 ng/mL –	8,000 units per day
30–40 ng/mL –	5,000 units per day
40–50 ng/mL –	2,000 units per day

If you are taking a vitamin D supplement, adequate calcium and magnesium intake are also required.

It is very difficult to get too much vitamin D. People can take up to 10,000 units per day for 6 months and not have any adverse effects. However, people with sarcoidosis, tuberculosis, Lyme disease, lymphoma, or kidney disease have to be supplemented carefully because of an increased risk of their blood calcium level becoming too high.

Rechecking Your Vitamin D Level

It is recommended that you have your vitamin D level rechecked within 2 weeks to 2 months after starting supplementation, depending on your medical and health condition. Other lab tests for calcium, ionized calcium, magnesium, and parathyroid hormone levels (PTH) may also be done during the recheck.

-
- ⁱ Dietary Supplement Fact Sheet: Vitamin D. *Office of Dietary Supplements (ODS)*. National Institutes of Health (NIH). Retrieved 2010-04 11.
- ⁱⁱ Cedric F. Garland, Dr PH, FACE, Edward D. Gorham, MPH, PhD, Sharif B. Mohr, MPH, Frank C. Garland, PhD. Vitamin D for cancer prevention: Global perspective. *Annals of Epidemiology*. Volume 19, Issue 7, Pages 468-483 (July 2009).
- ⁱⁱⁱ P. Lips, D. Hosking, K. Lippuner, J. M. Norquist, L. Wehren, G. Maalouf, S. Ragi-Eis, J. Chandler. The prevalence of vitamin D inadequacy amongst women with osteoporosis: an international epidemiological investigation. Volume 260, Issue 3, pages 245–254, September 2006
- ^{iv} Siegfried Segaer. Vitamin D regulation of cathelicidin in the skin: Toward a renaissance of vitamin D in Dermatology? *Journal of Investigative Dermatology* (2008) 128, 773–775. doi:10.1038/jid.2008.35
- ^v Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc*. 2003;78(12):1463-1470.
- ^{vi} Al Faraj, Saud MD; Al Mutairi, Khalaf MD. Vitamin D deficiency and chronic low back pain in Saudi Arabia. *Spine*:15 January 2003 - Volume 28 - Issue 2 - pp 177-179.
- ^{vii} Armstrong, D.; Meenagh, G.; Bickle, I.; Lee, A.; Curran, E.; Finch, M Vitamin D deficiency is associated with anxiety and depression in fibromyalgia. *Clinical Rheumatology*. Volume 26, Number 4, 551-554, DOI: 10.1007/s10067-006-0348-5
- ^{viii} Mathieu C, Gysemans C, Giuliatti A, Bouillon R. Vitamin D and diabetes. *Diabetologia*. 2006 Jan;49(1):217-8.
- ^{ix} Mackay-Sim A, Féron F, Eyles D, Burne T, McGrath J. Schizophrenia, vitamin D, and brain development. *Int Rev Neurobiol*. 2004;59:351-80.
- ^x Vitamin D Deficiency Common in Patients with Chronic Migraine. <http://www.medscape.com/viewarticle/577151>
- ^{xi} Ginanjar E, Sumariyono, Setiati S, Setiyohadi B. Vitamin D and autoimmune disease. *Acta Med Indones*. 2007 Jul-Sep;39(3):133-41.
- ^{xii} Lisa M. Bodnar, Janet M. Catov, Hyagriv N. Simhan, Michael F. Holick, Robert W. Powers and James M. Roberts. Maternal vitamin D deficiency increases the risk of preeclampsia. *The Journal of Clinical Endocrinology & Metabolism* Vol. 92, No. 9 3517-3522
- ^{xiii} *Int J Epidemiol*. 1990 Sep;19(3):559-63. Myocardial infarction is inversely associated with plasma 25-hydroxyvitamin D3 levels: a community-based study. Scragg R, Jackson R, Holdaway IM, Lim T, Beaglehole R. Department of Community Health, University of Auckland , New Zealand.
- ^{xiv} Li YC, Kong J, Wei M, Chen ZF, Liu SQ, Cao LP. 1,25-Dihydroxyvitamin D(3) is a negative endocrine regulator of the renin-angiotensin system. *J Clin Invest*. 2002;110(2):229-238.
- ^{xv} Ramagopalan SV, Maugeri NJ, Handunnetthi L, Lincoln MR, Orton S-M, et al. (2009) Expression of the multiple sclerosis-associated MHC Class II Allele *HLA-DRB1*1501* Is regulated by vitamin D. *PLoS Genet* 5(2): e1000369. doi:10.1371/journal.pgen.1000369
- ^{xvi} Anderson L, Cotterchio M, Vieth R, Knight J. Vitamin D and calcium intakes and breast cancer risk in pre- and postmenopausal women. *Am J Clin Nutr* 2010; 91(6): 1699-1701.
- ^{xvii} Garland CF, Gorham ED, Mohr SB, et al. Vitamin D and prevention of breast cancer: Pooled analysis. *J Steroid Biochem Mol Biol* 2007;103:708–11.
- ^{xviii} Bruce W Hollis and Carol L Wagner. Assessment of dietary vitamin D requirements during pregnancy and lactation. *American Journal of Clinical Nutrition*, Vol. 79, No. 5, 717-726, May 2004