





"Fueling the Passion"

2015 Concussion Management Program

Michael Stuart MD Alan Ashare MD

The standard of care for current medical practice and the law in most states requires that <u>any</u> <u>athlete with a suspected concussion</u> is **immediately removed from play**.

A concussion is a traumatic brain injury- there is no such thing as a minor brain injury.

A player does not have to be "knocked-out" to have a concussion-less than 10% of players actually lose consciousness.

A concussion can result from a blow to head, neck or body.

Concussions often occur to players who don't have or just released the puck, from open-ice hits, unanticipated hits and illegal collisions.

The youth hockey player's brain is more susceptible to concussion.

In addition, the concussion in a young athlete may be harder to diagnosis, takes longer to recover, is more likely to have a recurrence and can be associated with serious long-term effects.

Treatment is individualized and it is impossible to predict when the athlete will be allowed to return to play- *there is no standard timetable*.

A player with any symptoms or signs; disorientation; impaired memory, concentration, balance or recall has a concussion.

Remember these steps:

Remove immediately from play (training, practice or game)

Inform the player's coach/parents

Refer the athlete to a qualified health-care professional

Initial treatment requires physical and cognitive rest

The athlete begins a graded exertion and schoolwork protocol.

Medical clearance is required for return to play

Diagnosis

Players, coaches, parents and heath care providers should be able to recognize the symptoms and signs of a concussion:

Symptoms

Headache

Nausea

Poor balance

Dizziness

Double vision

Blurred vision

Poor concentration

Impaired memory

Light Sensitivity

Noise Sensitivity

Sluggish

Foggy

Groggy

Confusion

<u>Signs</u>

Appears dazed or stunned

Confused about assignment

Moves clumsily

Answers slowly

Behavior or personality changes

Unsure of score or opponent

Can't recall events after the injury

Can't recall events before the injury

Management Protocol

- 1. If the player is unresponsive- call for help & dial 911
- **2.** If the athlete is *not breathing*: start CPR

DO NOT move the athlete

DO NOT remove the helmet

DO NOT rush the evaluation

3. Assume a neck injury *until proven otherwise*

DO NOT have the athlete sit up or skate off until you have determined:

no neck pain

no pain, numbness or tingling

no midline neck tenderness normal muscle strength normal sensation to light touch

4. If the athlete is conscious & responsive without symptoms or signs of a neck injury...

help the player off the ice to the locker room perform an evaluation do not leave them alone

5. Evaluate the player in the locker room:

Ask about concussion *symptoms* (How do you feel?)

Examine for signs

Verify *orientation* (What day is it?, What is the score?, Who are we playing?)

Check *immediate memory* (Repeat a list of 5 words)

Test *concentration* (List the months in reverse order)

Test *balance* (have the players stand on both legs, one leg and one foot in front of the other with their eyes closed for 20 seconds)

Check *delayed recall* (repeat the previous 5 words after 5-10 minutes)

6. A player with any symptoms or signs, disorientation, impaired memory, concentration, balance or recall has a concussion

"When in doubt, sit them out"

Remove immediately from play (training, practice or game Inform the player's coach/parents
Refer the athlete to a qualified health-care professional
Medical clearance is required for return to play

7. If any of the signs or symptoms listed below develop or worsen: go to the hospital emergency department or dial 911.

Severe throbbing headache

Dizziness or loss of coordination

Memory loss or confusion

Ringing in the ears (tinnitus)

Blurred or double vision

Unequal pupil size

No pupil reaction to light

Nausea and/or vomiting

Slurred speech

Convulsions or tremors

Sleepiness or grogginess

Clear fluid running from the nose and/or ears

Numbness or paralysis (partial or complete)

Difficulty in being aroused

8. An athlete who is *symptomatic* after a concussion initially requires *physical* and *cognitive rest*.

A concussed athlete should not participate in physical activity, return to school, play video games or text message if he or she is having symptoms at rest.

Concussion symptoms & signs *evolve over time*- the severity of the injury and estimated time to return to play are unpredictable.

9. A qualified health care provider guides the athlete through a **Functional Return to School/Play protocol**.

USA Hockey Post-Concussion Functional Return to School/Play Protocol

This protocol should not be initiated until after the athlete has been released to participate in the functional return to play protocol by a qualified health care provider. *If symptoms worsen during a functional test, the test should be stopped and the athlete monitored until symptoms resolve.* No further functional testing should be performed that day. Functional testing may resume the following day at the previously asymptomatic level if the athlete remains asymptomatic. If symptoms do not resolve, appropriate medical attention should be obtained.

After each phase of functional testing, the presence of post-concussive symptoms should be assessed and progression to the next phase of functional testing will require the absence of post-concussive symptoms. Each phase requires *a minimum of 1 day* before progressing to the next phase.

Level 2: Light aerobic exercise such as walking or stationary cycling. No resistance training. May read 20-40% of normal volume. Limited TV, but No video games. Partial school day as tolerated.

Level 3: Sport-specific exercises. No resistance training. May read 40-60 % of normal volume. No video games. Full school day as tolerated.

Level 4: Non-contact practice. Resistance training OK. No video games. Full school day

Level 5: Full contact practice. Full school day. Video games OK.

Level 6: Return to unrestricted competition and cognitive activities if medical clearance is provided by a qualified health care provider